

DNA Barcoding of The Soft Coral, *Clavularia inflata*, Shows Two Major Groups Across Indonesian Coral Reefs

Beginer Subhan^{1,*}, Dietrich G. Bengen¹, Sebastian Ferse^{2,3}, Fauzan Dzulfannazhir¹,
Luzmi Malia Izza¹, Nurlita Putri Anggraini¹, Prakas Santoso¹, Dondy Arafat¹, Lalu M. Iqbal Sani⁴,
Hawis Madduppa^{1,4}

¹Department of Marine Science and Technology, Faculty of Fisheries and Marine Science, IPB University
Kampus IPB Dramaga Bogor 16680 West Java, Indonesia

²Leibniz Centre for Tropical Marine Research,
Fahrenheitstraße 6, 28359 Bremen, Germany

³Department of Marine Ecology, Faculty of Biology and Chemistry, University of Bremen
Bibliothekstraße 1, 28359 Bremen, Germany

⁴Oceanogen Environmental Biotechnology Laboklinikum
Jl. Bambu Ori 3 No. 46, Taman Yasmin Sektor 7, East Cilendek, Bogor, West Java 16112, Indonesia
Email: beginersubhan@apps.ipb.ac.id

Abstract

Clavularia inflata was first described from Ternate Island, Indonesia in 1896 and later reported ~~appeared~~ from Japan and Taiwan in 1953. *Clavularia* (Blainville 1830) soft corals exhibit complex morphological traits that are difficult to differentiate, thus complicating their identification. DNA barcoding has been envisioned and actively pushed as a credible method for assigning unidentified specimens to known species by comparison to a molecular reference data database. DNA barcoding is an important tool to support the management of biological resources, one of which are soft corals. As soft corals are among the species that are difficult to identify morphologically, DNA barcoding data are very valuable for identifying them. Thus, the purpose of this study was to use molecular methods to confirm the identity of 25 colonies taken from 13 Indonesian coral reef sites (Anambas, Natuna, Tanjung Lesung, Kepulauan Seribu, Madura, Kangean, Bontang, Lombok, Manado, Wakatobi, Morotai, Kon, and Maluku Tenggara Barat (MTB)) and putatively identified as *C. inflata*. All specimens were identified as *C. inflata* molecularly using the mitochondrial DNA *mtMuts* gene. The results from this study can enrich the information base for barcoding of soft corals elsewhere because the samples were taken across a range of nearly 4000km throughout Indonesia archipelago. Although a comparison of the nucleotide base chains to Genbank data indicates that the samples belong to a single species, two clades in the phylogenetic tree and data from the Automatic Barcode Gap Discovery (ABGD) indicate that there are two major groups of *C. inflata* in Indonesia, implying cryptic species.

Keywords: Molecular taxonomy, Phylogeny, Octocorallia, DNA Barcoding

Introduction

The sub-class Octocorallia (Cnidaria: Anthozoa) contains more than 2,000 species which are spread all over the world and occur in a variety of habitats (Bayer, 1973; Fabricius and Alderslade, 2001). Octocorals have polyps with eight pinnate tentacles and internal calcareous skeletal elements, termed sclerites. The octocoral order Alcyonacea (soft corals) is the largest in number of species. The species of this order may or may not contain symbiotic algae in their tissue (they are zooxanthellate vs. azooxanthellate species, respectively). Alcyonacea are found in abundance in coral reef habitats in most of the Indo-West Pacific and play an important ecological role, often occurring in percentages of benthic substrate

cover equal to or greater than that of scleractinian corals (Tursch and Tursch, 1982; Dinesen, 1983; Dai, 1988; Riegl et al., 1995; Fabricius, 1997).

There are about 96 soft coral genera belonging to 23 families in the Indo-Pacific (Fabricius and Alderslade, 2001). Manuputty (2002) notes 28 genera of soft corals in 4 families occurring in Indonesian waters, one of which is *Clavularia*. *Clavularia* (Blainville, 1830) spp., belonging to the suborder stolonifera, are zooxanthellate soft corals, with every single polyp connected to the base. Polyps can be pink to brownish gray and measure about 30 mm in diameter. Stolons are brown to dark red, and are often covered by algae or sponges (Fabricius and Alderslade, 2001). *Clavularia inflata* was first described from Ternate island, Indonesia (Schenk,

1896). Utinomi (1953) later reported *C. inflata* from Japan and Taiwan.

DNA barcoding has been envisioned and widely promoted as a reliable technique by which unidentified specimens can be assigned to known species by comparison to a reference database of molecular exemplars (Hebert *et al.*, 2003). At this time the use of DNA barcodes is widely spread because it has several advantages: (1) DNA accurately distinguishes groups that have the same morphological character (Bingpeng *et al.*, 2018), (2) it can identify species at different stages of the developmental cycle (Hubert *et al.*, 2010), and (3) it can be used to help identify specimens that are damaged and incomplete (Prehadi *et al.*, 2015). Similarly, it can be used to identify material such as processed fish or shellfish found in seafood products and thus assist in the authentication of fishery products (Abdullah *et al.*, 2020) for traceability of marine fisheries. A molecular approach has been used successfully to identify marine biota at the species level in a number of different taxa, *i.e.* soft corals (McFadden *et al.*, 2006a; McFadden *et al.*, 2009; Benahayu and McFadden 2011; Mc Fadden *et al.*, 2014; Kusuma *et al.*, 2016), fish (Jefri *et al.*, 2015; Prehadi *et al.*, 2015; Sembiring *et al.*, 2015; Madduppa *et al.*, 2016; Toha *et al.*, 2020; Fadli *et al.*, 2020; Madduppa *et al.*, 2021), giant clams (DeBoer *et al.*, 2008), sea squirts (Anzani *et al.*, 2019), coral reef fish (Ackiss *et al.*, 2013), small crustaceans (Hazeri *et al.*, 2019), sea turtles (Madduppa *et al.*, 2019), sponges (Madduppa *et al.*, 2015), horse shoe crabs (Aini *et al.*, 2020), crown-of-thorns starfish (Vogler *et al.*, 2013), cephalopods (Kholilah *et al.*, 2021), gastropods (Selaky *et al.*, 2017), and sea cucumbers (Madduppa *et al.*, 2017).

Soft coral research in Indonesia has been limited to assessing the biological aspects and distribution of soft corals (Manuputty, 2002, 2008, 2010). Phylogenetic relationships in *Clavularia* tropical soft corals have been studied by McFadden and Ofwegen (2012). To date, DNA barcoding has been carried out for a number of species of Clavulariidae: *Clavularia borealis* (Moore *et al.*, 2017), *Hanabira yukibana* (Lau *et al.*, 2019), *Inconstantia procera*, *I. pannucea*, *Carijoa riisei*, *Arula petunia*, *Cornularia pabloi* (Mc Fadden and Ofwegen, 2012) and *Trachythela rudis* (Quattrini *et al.*, 2014).

More recently, the use of molecular identification for soft corals has also become a supporting tool in describing or determining new species (Benahayu and McFadden, 2011). The results of identification can also aid in delineating boundaries between species that have not been revealed previously (McFadden *et al.*, 2006, 2009; France, 2007). The genetic information for marine

organisms in Indonesia is still sparse (Madduppa *et al.*, 2021). Therefore, this study aimed to apply DNA barcoding on samples putatively identified as *Clavularia inflata* collected from a suite of Indonesian coral reefs in order to verify species identity and assess the potential existence of cryptic species.

Material and Methods

Study sites, specimen and tissue sampling

Sampling of specimens and tissue was conducted sporadically between February and August 2018 at 13 sites (Anambas, Natuna, Tanjung Lesung, Kepulauan Seribu, Madura, Kangean, Bontang, Lombok, Manado, Wakatobi, Morotai, Kon, and Maluku Tenggara Barat; Table 1.).

Sampling was carried out using a purposive sampling approach. Purposive sampling is a nonprobability sampling technique in which the researcher selects the sample according to its presence (Etikan *et al.*, 2016). This can reduce the risk of an insufficient number of samples. Furthermore, the selection of research sites considered the spread of *Clavularia inflata* in Indonesia and information from diving clubs and divers. Samples were taken from between 4 and 7 colonies at each site. A colony, referring to an individual, was defined as a discrete patch of polyps connected by stolons, and only colonies more than 3 m apart were sampled (Bastidas *et al.*, 2002). A total of 25 sample mtDNA sequences were obtained from 13 locations across Indonesia coral reefs (Figure 1).

During the dive, each sample was stored in a sample tube filled with seawater. Back on board, samples were transferred into sterile sample bottles of 10 ml volume filled with 97% ethanol. Upon arrival in the laboratory, the samples were then stored in the freezer at a temperature of 4 °C. Sampled colonies were documented before each sample was taken using a Canon G16 camera in underwater housing. For each colony, a wide-angle photo of the colony and its surrounding environment and a close-up photo were taken.

Molecular analysis

DNA was extracted from tissue samples using the gSYNC DNA Extraction Kit for Blood and Tissue from Geneaid. DNA extraction followed the standard protocol of the manufacturer, modified and optimized for the genomic DNA extraction from *Clavularia inflata*. Approximately 40 mg of tissue sample were ground and put into 1.5 ml microcentrifuge tubes, 150 µl GST Buffer and 20 µl Proteinase K were added, and the samples were then vortexed. The samples were incubated at 60 °C over night.




Figure 1. Sites of tissue and specimen sampling at 13 coral reef locations across Indonesia

Table 1. Sampling sites and number of samples of *Clavularia inflata* across Indonesian coral reefs

Sampling sites	Number of samples	Position
Anambas	2	3°23'27.2"N 106°08'56.0"E
Natuna	2	4°00'12.1"N 108°24'45.8"E
Tanjung Lesung	2	6°28'43.7"S 105°39'31.3"E
Kepulauan Seribu	2	5°38'38.4"S 106°33'25.0"E
Madura	1	7°12'03.7"S 114°02'55.8"E
Kangean	2	6°50'33.7"S 115°13'40.1"E
Lombok	2	8°43'25.7"S 115°57'42.5"E
Bontang	2	0°04'40.6"N 117°33'50.9"E
Manado	3	1°32'42.9"N 124°48'55.2"E
Wakatobi	1	5°28'05.9"S 123°42'49.2"E
Morotai	2	2°02'42.1"N 128°17'27.4"E
Kon	2	3°55'26.8"S 131°12'33.8"E
Maluku Tenggara Barat (MTB)	2	8°14'41.4"S 130°42'28.3"E

Mitochondrial DNA mtMuts partial gene fragments were amplified by using ND42625F (5-TACGTGGYACAATTGCTG-3) (Lepard 2003) and Mut-3458R (5-TSGAGCAAAGCCACTCC-3) (Sanchez *et al.*, 2003). The PCR reaction was carried out in a total volume of 25 µl, including 12.5 µl MyTaq™ Red Mix – Bionline, 10 mM of each primer (Forward and Reverse) at a volume of 1.25 µl, 9 µl deionized water (ddH₂O), and 1 µl DNA template. The mixture was run in a thermal cycler by using the following PCR cycle: pre-denaturation at 95 °C for 3 mins, followed by 35 cycles of denaturation at 95 °C for 30 sec, annealing at 50 °C for 30 sec, and extension at 72 °C for 1 min.

DNA product quality was checked by using an electrophoresis chamber. The 1 % agarose gel and DNA dye Ethidium Bromide (4µl) were used as gel agarose media of electrophoresis. A total of 1µl PCR product was loaded and inserted in agarose wells. The electrophoresis process was run using an

electrophoresis machine with a voltage of 100 V-135V and a current of 400 mA, with a time of 35 minutes. Electrophoresis results were visually inspected using an Alphamager Mini doc gel machine, and the photos of the results stored digitally. PCR products of sufficient quality were then sequenced by using a modified Sanger protocol (Sanger *et al.*, 1977) at Firstbase Company, Malaysia.

Data analysis

Sequences were edited and aligned using MEGA 6.06 (Molecular Evolutionary Genetic Analysis) (Tamura *et al.*, 2013). After editing and aligning, DNA sequences were trimmed to reduce the negative impact from alien oligonucleotide sequences in the read end (Criscuolo and Brisse, 2013). The aligned data was matched in the NCBI (National Center for Biotechnology Information) search system using BLAST (Basic Local Alignment Search Tool) accessed

via the website <http://blast.ncbi.nlm.nih.gov>, and percent identity and query cover noted. The percent identity is a quantity that explains how similar the query sequence is to the target sequence (number of characters in each sequence that are identical). The higher the percent identity is, the more significant the match. The query cover is a number that describes how much of the query sequence is covered by the target sequence. All sequences were deposited in BOLD systems under the project “DNA barcoding of Clavulariidae in Indonesia” (<http://www.boldsystems.org>). Species delineation by determining the number of Operational Taxonomic Units (OTUs) based on pairwise sequence distances between individuals within the dataset was applied with the Automatic Barcode Gap Discovery (ABGD) tool (Puillandre *et al.*, 2012).

Phylogenetic tree reconstruction with Neighbor-Joining and Maximum Likelihood was performed in MEGA 6.06. Re-sampling characteristics used non-parametric bootstrapping with 1000 replications per bootstrap to visualize the branch in

MEGA 6.06. For DNA evolution, there are many models that can be used to reconstruct a phylogenetic relationship. Initially, the best-fitting model was searched by using Find Best DNA/Protein (ML) in the MEGA 6.06 Program. The Tamura 3-parameter model was used to select the best-fitting model based on maximum likelihood. To analyze genetic distance, the Neighbor-Joining (NJ) method and a Kimura 2-parameter model were used. 1000 replicates per bootstrap were used, and out-grouped with the species *Eunicea flexuosa* (accession number: KC310539.1) to root the trees.

Results and Discussion

The molecular identification (Table 2-) confirmed that all tissue samples belonged to *Clavularia inflata*, with a high percent identity (92% - 100%) and query cover of 95-100%. Most of the percent identity was between 99-100%; only four of the 25 samples (from Kon and Bontang) had a percent identity value below 99%.

Table 2. Quality of molecular identification of sample sequences collected from the different sampling sites, showing ID numbers used in BLAST searches on NCBI, maximum identification (%), and query cover (%). All samples were identified as belonging to *Clavularia inflata*.

ID Number	Sampling sites	Max. Ident (%)	Query (%)	Cover
ITK SC ANAMBAS 02	Anambas	99		99.84
ITK SC ANAMBAS 01	Anambas	99		100
ITKSCNAT17	Natuna	100		100
ITKSCNAT13	Natuna	100		100
ITKTLSC23	Tanjung Lesung	99		100
ITKTLSC09	Tanjung Lesung	99		100
ITKPJGSC10	Kep Seribu	99		100
ITKPJGSC09	Kep Seribu	99		100
ITK MDR SC 05	Madura	99		100
KANG 01	Kangean	99		100
KANG 02	Kangean	99		100
ITK LMBK SC 1_12	Lombok	99		100
ITK LMBK SC 1_06	Lombok	99		100
ITK BON SC 32	Bontang	94		99.46
ITK BON SC 35	Bontang	94		99.46
ITK MER SC 03	Manado	100		99.49
ITK MER SC 02	Manado	100		99.49
ITK WKT SC 09	Wakatobi	99		95.76
ITK MER SC 10	Manado	100		99.49
ITK MOR SC 05	Morotai	100		100
ITK MOR SC 07	Morotai	99		100
ITK KON SC 07	Kon	93		99.82
ITK SC KON 16	Kon	92		100
ITK MTB SC 48	Maluku Tenggara Barat	99		99.96
ITK MTB SC 50	Maluku Tenggara Barat	99		99.96

Table 3. Summary statistics for nucleotide frequency distribution of *Clavularia inflata* sequences (n = 25) across the 13 sampled Indonesian coral reef locations

Nucleotide base	Min	Mean	Max
C%	16.24	16.35	17.29
G%	18.78	19.31	20.47
A%	28.60	30.22	30.63
T%	33.28	34.12	34.52




Figure 2. Phylogenetic tree of 25 sequences *Clavularia inflata* constructed using (A) Neighbor Joining (NJ) and (B) the Maximum Likelihood method with Kimura 2 Parameter model, bootstrap value 1000x.

The results of genetic analysis confirmed the entire study samples as belonging to *Clavularia inflata*. The phylogenetic trees constructed from the 25 nucleotide base chain sequences derived from the samples using Neighbor Joining and Maximum Likelihood show two large clades (Figure 2.).

The 25 sequences generated from *Clavularia inflata* show a nucleotide composition of A= 30.22%, T= 34.12%, C= 16.35%, G= 19.31% (Table 3.). The nucleotide composition of all sequences was related to the genetic distances between individuals. Sample ITK WKT SC 09 from Wakatobi had the highest genetic distance to other samples in this study (Table 4.). The number of OTUs generated by ABGD based on the Kimura-2-parameter model (K2P) varied from

2 to 5 (Figure 3.). The initial partition at a prior intraspecific divergence (P)= 0.001–0.007 produced 3 OTUs, while a Recursive partition at a prior intraspecific divergence (P)= 0.0129–0.1 produced 2 OTUs.

All samples used in this study could be unequivocally identified as belonging to *C. inflata* by means of molecular tools. The use of molecular identification for soft corals has become a regular tool for describing or determining new species (Benahayu and McFadden, 2011). Furthermore, the results of molecular identification can also provide support in delineating boundaries between species that have not been previously revealed (France, 2007; McFadden et al., 2009).

DNA sequences of *C. inflata* are still rarely found in genetic databases. For example, there are only eight mtMuts sequences in GenBank, each originating from either Okinawa/Japan or Palau. As this study has contributed an additional 25 *Clavularia inflata* sequences to a database that is freely available to the public, the results from this study can enrich the information base for barcoding of soft corals elsewhere because the samples were taken across a range of nearly 4000km throughout Indonesia. The available sequences will also be beneficial for DNA barcoding identification. Additional information from this research can help in studying and deciphering complexities in the clavulariidae family in particular or soft corals in general.

Information from ABGD shows that there are two major groups of *C. inflata* found in Indonesia. The two groups were also seen in the phylogenetic tree, namely the group from the first clade originating from Wakatobi and Manado and the second clade from samples originating from Tanjung Lesung, Kepulauan Seribu, Maluku Tenggara Barat, Lombok, Madura, Anambas, Bontang, Kon, Natuna, Kangean and Morotai. Currents are one of the factors that causes genetic similarity in the same species in different locations (Knittweis *et al.*, 2009). Wakatobi and Manado are passed by the ITF which moves through the Pacific, passing by Manado and then to the south via Wakatobi. This is likely responsible for the higher genetic closeness at these compared to other locations. The observation of two major groups indicates the possibility of the presence of cryptic

species in this type. The potential for identification of cryptic species and genera using mtDNA barcodes is still high, as evidenced by the outcome of biodiversity studies in the Red Sea and elsewhere (McFadden *et al.*, 2006a, 2009). Previous research using visual tools identified *C. inflata* var. *luzoniana* May 1899 (Roxas, 1933) in the Philippines, which has different morphological characteristics from *Clavularia inflata*. However, genetic information about the variant is not yet available, so this study cannot be compared genetically. The hypothetical existence of two species in this study region could eventually be confirmed by using other primers on different loci of DNA. The species partition hypothesis needs further exploration, such as the inclusion of more than one locus as additional information, since a single locus is not strong enough to develop reliable species hypotheses (Puillandre *et al.*, 2012). In a previous study of *Alcyonium*, interspecific genetic distances were twice greater for *msh1* than COI, each marker identified similar numbers of species, and the extended COI + *igr1* + *msh1* barcode more effectively discriminated sister taxa (McFadden *et al.*, 2011).

DNA barcoding is an important tool to support the management of biological resources, one of which are soft corals. As soft corals are among the species that are difficult to identify morphologically, DNA barcoding data are very valuable for identifying them. Studies that combine molecular character-state data (*i.e.* barcodes) with traditional morphological taxonomic approaches have greatly improved our understanding of species boundaries




Figure 4. The number of genetically distinct Operational Taxonomic Units (OTUs) according to the recursive partition (red dots) and initial partition (yellow dots) at a prior intraspecific divergence value generated by Automatic Barcode Gap Discovery (ABGD) based on the Kimura-2-parameter model (K2P).

Table 4. Pairwise comparisons based on mean Kimura-2-parameter (K2P) distances between *Clavularia inflata* across the sampled Indonesian coral reef locations. Locations for each sample ID are given in Table 2

Sample Id	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	
ITK SC ANAMBAS 02																										
ITK SC ANAMBAS 01	0,003																									
ITK BON SC 32	0,038	0,038																								
ITK BON SC 35	0,038	0,038	0,003																							
KANG 01	0,002	0,002	0,04	0,04																						
KANG 02	0,002	0,002	0,04	0,04	0																					
ITK KON SC 07	0,03	0,033	0,031	0,031	0,031	0,031																				
ITK SC KON 16	0,042	0,042	0,04	0,04	0,044	0,044	0,03																			
ITK LMBK SC 12	0,002	0,002	0,04	0,04	0	0	0,031	0,044																		
ITK LMBK SC 06	0,002	0,002	0,04	0,04	0	0	0,031	0,044	0																	
ITK MDR SC 05	0,002	0,002	0,04	0,04	0	0	0,031	0,044	0	0																
ITK MER SC 03	0,012	0,009	0,033	0,033	0,01	0,01	0,042	0,047	0,01	0,01	0,01															
ITK MER SC 02	0,012	0,009	0,033	0,033	0,01	0,01	0,042	0,047	0,01	0,01	0,01	0														
ITK MOR SC 05	0,007	0,003	0,038	0,038	0,005	0,005	0,037	0,042	0,005	0,005	0,005	0,005	0,005													
ITK MOR SC 07	0,003	0	0,038	0,038	0,002	0,002	0,033	0,042	0,002	0,002	0,002	0,009	0,009	0,003												
ITK MTB SC 48	0,005	0,005	0,044	0,044	0,003	0,003	0,035	0,047	0,003	0,003	0,003	0,014	0,014	0,009	0,005											
ITK MTB SC 50	0,005	0,005	0,044	0,044	0,003	0,003	0,035	0,047	0,003	0,003	0,003	0,014	0,014	0,009	0,005	0										
ITK SCNAT17	0,007	0,003	0,038	0,038	0,005	0,005	0,037	0,042	0,005	0,005	0,005	0,005	0,005	0	0,003	0,009	0,009									
ITK SCNAT13	0,007	0,003	0,038	0,038	0,005	0,005	0,037	0,042	0,005	0,005	0,005	0,005	0,005	0	0,003	0,009	0,009	0								
ITK PJGSC10	0,002	0,002	0,04	0,04	0	0	0,031	0,044	0	0	0	0,01	0,01	0,005	0,002	0,003	0,003	0,005	0,005							
ITK PJGSC09	0,002	0,002	0,04	0,04	0	0	0,031	0,044	0	0	0	0,01	0,01	0,005	0,002	0,003	0,003	0,005	0,005	0						
ITK TLSC23	0,002	0,002	0,04	0,04	0	0	0,031	0,044	0	0	0	0,01	0,01	0,005	0,002	0,003	0,003	0,005	0,005	0	0					
ITK TLSC09	0,005	0,002	0,04	0,04	0,003	0,003	0,035	0,044	0,003	0,003	0,003	0,007	0,007	0,002	0,002	0,007	0,007	0,002	0,002	0,003	0,003	0,003				
ITK WKT SC 09	0,048	0,044	0,066	0,066	0,046	0,046	0,073	0,073	0,046	0,046	0,046	0,038	0,038	0,044	0,044	0,049	0,049	0,044	0,044	0,046	0,046	0,046	0,046			
ITK MER SC 10	0,012	0,009	0,033	0,033	0,01	0,01	0,042	0,047	0,01	0,01	0,01	0	0	0,005	0,009	0,014	0,014	0,005	0,005	0,01	0,01	0,01	0,007	0,038		

and intracellular relationships and have identified new, diagnostic morphological character states useful for field identification in several speciose and ecologically important genera of octocorals (McFadden *et al.*, 2006a, 2009; France, 2007). *C. inflata* is known to possess biomedical properties (Sheu *et al.*, 2019) but is also traded as an ornamental animal for marine aquaria (Wabnitz *et al.*, 2003). The data presented here can later be used as a basis for tracking the sources of invasive origin or introduced species in other regions. For example, observations in Brazil have found *Clavularia* cf. *viridis* soft corals invading benthic substrate. The presence of these invertebrates is assumed to originate from the ornamental aquarium trade sourced from the Indo-Pacific region (Mantelatto *et al.*, 2018).

Conclusion

Molecular analysis confirmed all samples as belonging to *C. inflata*. The combination of different genetic tools in this study suggests the presence of cryptic species in this type. Further studies using additional molecular markers and morphological characteristics are suggested to further explore the possibility of cryptic species.

Acknowledgement

This study was funded by the Ministry of Research, Technology and Higher Education of the Republic of Indonesia in 2018 under the scheme "Penelitian Disertasi Doctor (PDD)". This study was supported by the Marine Biodiversity and Biosystematics Laboratory, Department of Marine Science and Technology, IPB University, Bogor, West Java, Indonesia. This paper is supported by USAID through Sustainable Higher Education Research Alliances (SHERA) Program- Animal Biotechnology and Coral Reef Fisheries (ANBIOCORE). The authors would like to thank Miss Nurlita Putri Anggraeni for technical laboratory support.

References

- Abdullah, A., Nurilmala, M., Muttaqin, E. & Yulianto, I., 2020. DNA-based analysis of shark products sold on the Indonesian market towards seafood labelling accuracy program. *Biodiversitas*, 21(4): 1385-1390. <https://doi.org/10.13057/biodiv/d210416>
- Ackiss, A.S., Pardede, S., Crandall, E.D., Ablan-Lagman, M.C.A., Barber, P.H. Ambariyanto, A. & Carpenter, K.E., 2013. Pronounced genetic structure in a highly mobile coral reef fish, *Caesio cuning*, in the Coral Triangle. *Mar. Ecol. Prog. Ser.*, 480: 185-197.
- Aini, N.K., Erwyansyah, E., Kurnia, R., Butet, N.A., Mashar, A., Zairion, Z., Funch, P., Madduppa, H. & Wardiatno, Y. 2020. Two morphotypes of tri-spine horseshoe crab, *Tachypleus tridentatus* (Leach, 1819) (Xiphosura Limulidae) in Indonesia and implications for species identification. *Biodiversity*, 11(2): 535-540. <https://doi.org/10.31396/Biodiv.Jour.2020.11.2.535.540>
- Anzani L., Madduppa, H., Nurjaya, I.W. & Dias, P.J. 2019. Short communication: Molecular identification of white sea squirt *Didemnum* sp. (tunicata, ascidiacea) colonies growing over corals in Raja Ampat Islands, Indonesia. *Biodiversitas*. 20(3): 636-642. <https://doi.org/10.13057/biodiv/d200304>
- Bayer F.M. 1973. Colonial organization in octocorals. In: Brodman R.S., A.H. Cheetman & W.A. Oliver Jr. (Ed). Animal colonies development and function through time. Dowden, Hutchinson and Ross. Pennsylvania. P: 69-93
- Bastidas, C., Benzie, J. & Fabricius, K 2002. Genetic differentiation among populations of the brooding soft coral *Clavularia koellikeri* on the Great Barrier Reef. *Coral Reefs*, 21(3): 233-241. <https://doi.org/10.1007/s00338-002-0236-7>
- Benayahu Y. & McFadden, C.S. 2011. A new genus of soft coral of the family Alcyoniidae (Cnidaria, Octocorallia) with re-description of a new combination and description of a new species. *ZooKeys*. 84: 1-11. <https://doi.org/10.3897/zookeys.84.781>
- Bingpeng, X., Heshan, L., Zhilan, Z., Chunguang, W., Yanguo, W. & Jianjun, W. 2018. DNA barcoding for identification of fish species in the Taiwan Strait. *Plos One*. 13: e0198109. <https://doi.org/10.1371/journal.pone.0198109>
- Criscuolo A. & Brisse, S. 2013. Alien Trimmer: a tool to quickly and accurately trim off multiple short contaminant sequences from high-throughput sequencing reads. *Genomics*. 102(5-6): 500-506. <https://doi.org/10.1016/j.ygeno.2013.07.011>
- Dai, C.F. 1988. Coral communities of southern Taiwan. *Proc 6th Intl Coral Reef Symp.* 2: 647-652.
- DeBoer, T.S., Subia, M.D., Ambariyanto, A., Erdmann, M.V., Kovitvongsa, K. & Barber, P. H. 2008. Phylogeography and limited genetic connectivity in the endangered boring giant clam across the Coral Triangle. *Conserv. Biol.*, 22(5): 1255-1266.

- Dinesen, Z.D. 1983. Patterns in the distribution of soft corals across the central Great Barrier Reef. *Coral Reefs*. 1: 229-236. <https://doi.org/10.1007/BF00304420>
- Dray S. & Dufour, A.B. 2007. The ade4 package: Implementing the duality diagram for ecologists. *J. Stat. Softw.*, 22: 1-20. <https://doi.org/10.18637/jss.v022.i04>.
- Etikan I., Musa, S.A. & Alkassim, R.S. 2016. Comparison of convenience sampling and purposive sampling. *Am. J. Theor. Appl. Stat.*, 5(1): 1-4. doi: 10.11648/j.ajtas.20160501.11
- Fadli, N., Nor, S.A.M., Othman, A.S., Sofyan, H. & Muchlisin, Z.A. 2020. DNA barcoding of commercially important reef fishes in Weh Island, Aceh, Indonesia. *PeerJ*. 8: e9641 <https://doi.org/10.7717/peerj.9641>
- Fabricius K.E. 1997. Soft coral abundance on the central Great Barrier Reef: effects of *Acanthaster planci*, space availability, and aspects of the environment. *Coral Reefs*. 16: 159-167. <https://doi.org/10.1007/s003380050070>
- Fabricius K.E. & Alderslade, P. 2001. Soft corals and sea fans: a comprehensive guide to the tropical shallow water genera of the central-west Pacific, the Indian Ocean and the Red Sea. Australian Institute of Marine Science, Townsville.
- France, S.C. 2007. Genetic analysis of bamboo corals (Cnidaria: Octocorallia: Isididae): does lack of colony branching distinguish *Lepidisis* from *Keratoisis*. *Bull. Mar. Sci.*, 81: 323-333.
- Hazeri, G., Rahayu, D.L., Subhan, B., Sembiring, A., Anggoro, A.W., Ghazali, A.T. & Madduppa, H.H. 2019. Latitudinal species diversity and density of cryptic crustacean (brachyura and anomura) in micro-habitat autonomous reef monitoring structures across Kepulauan Seribu, Indonesia. *Biodiversitas*. 20(5): 1466-1474. <https://doi.org/10.13057/biodiv/d200540>
- Hebert, P.D., Cywinska, A., Ball, S.L. & DeWaard, J.R. 2003. Biological identifications through DNA barcodes. *Proc. Royal Soc. B*. 270: 313-321. <https://doi.org/10.1098/rspb.2002.2218>
- Hubert N., E. Delrieu-Trottin, J-O. Irisson, C. Meyer & S. Planes. 2010. Identifying coral reef fish larvae through DNA barcoding: a test case with the families Acanthuridae and Holocentridae. *Mol. Phylogenet. Evol.*, 55: 1195-1203. <https://doi.org/10.1016/j.ympev.2010.02.023>
- Hoeksema B.W. 2007. Delineation of the Indo-Malayan centre of maximum marine biodiversity In: Willem Renema (Eds.), Biogeography, Time, and Place: Distributions, Barriers, and Islands *Topics in Geobiology*. 29: 117-178. https://doi.org/10.1007/978-1-4020-6374-9_5
- Hosein, F.N., Austin, N., Maharaj, S., Johnson, W., Rostant, L., Ramdass, A.C. & Rampersad, S.N. 2017. Utility of DNA barcoding to identify rare endemic vascular plant species in Trinidad. *Ecol. Evol.*, 7(18): 7311-7333. <https://doi.org/10.1002/ece3.3220>.
- Jefri, E., Zamani, N.P., Subhan, B. & Madduppa, H.H. 2015. Molecular phylogeny inferred from mitochondrial DNA of the grouper *Epinephelus* spp. in Indonesia collected from the local fish market. *Biodiversitas*. 16: 254-263. <https://doi.org/10.13057/biodiv/d160221>
- Kholilah N., Afiati, N. & Subagiyo. 2021. Characterization of Some Commercially Important Octopus (Mollusca: Cephalopoda) from Indonesian Waters using Mitochondrial DNA Cytochrome Oxidase SubUnit I (MtDNA COI). *Ilmu Kelautan: Indonesia Journal of Marine Sciences*. 26: 17-26. <https://doi.org/10.14710/ik.ijms.26.1.17-26>
- Kusuma A.B., Bengen, D.G., Madduppa, H., Subhan, B. & Arafat, D. 2016. Genetic diversity of soft coral *Sarcophyton trocheliophorum* in Java, Nusa Tenggara and Sulawesi Sea populations. *J. Enggano*. 1(1): 89-96. <https://doi.org/10.31186/jenggano.1.1.89-96>
- Knittweis, L., Kraemer, W.E., Timm, J. & Kochzius, M., 2009 Genetic structure of *Heliofungia actiniformis* (Scleractinia: Fungiidae) populations in the Indo-Malay Archipelago: implications for live coral trade management efforts. *Conserv. Genet.* 10(1): 241-249. <https://doi.org/10.1007/s10592-008-9566-5>
- Lau Y.W., Stokvis, F.R., Imahara, Y. & Reimer, J.D. 2019. The stoloniferous octocoral, *Hanabira yukibana*, gen. nov., sp. nov., of the southern Ryukyus has morphological and symbiont variation. *Contributions to Zoology*. 88(1): 54-77. <https://doi.org/10.1163/18759866-20191355>
- Lepard, A.B. 2003. Analysis of variation in the mitochondrial encoded MSH1 in the genus *Leptogorgia* (Cnidaria: Octocorallia) and implications for population and systematics studies. M.S. thesis, University of Charleston, Charleston, SC.

- Madduppa H., Cahyani, N.K.D., Anggoro, A.W., Subhan, B., Jefri, E., Sani, L.M.I., Arafat, D., Akbar, N. & Bengen, D.G. 2021. eDNA metabarcoding illuminates species diversity and composition of three phyla (chordata, mollusca and echinodermata) across Indonesian coral reefs. *Biodivers. Conserv.* 30(11): 3087-3114. <https://doi.org/10.1007/s10531-021-02237-0>
- Madduppa H., Putri A.S., Wicaksono R.Z., Subhan B., Akbar N., Ismail F., Arafat D., Prabuning D., Sani L.M.I., Srimariana E.S., Baksir A. 2020. Morphometric and DNA Barcoding of endemic Halmaheran walking shark (*Hemiscyllium halmahera*, Allen, 2013) in North Maluku, Indonesia: Morphogenetic of endemic Halmaheran walking shark. *Biodiversitas*, 21(7): 3331-3343. <https://doi.org/10.13057/biodiv/d210757>
- Madduppa, H., Subhan, B., Anggraini, N.P., Fadillah, R. & Tarman, K. 2017. Short communication: Dna barcoding reveals vulnerable and not evaluated species of sea cucumbers (Holothuroidea and Stichopodidae) from Kepulauan Seribu reefs, Indonesia. *Biodiversitas*. 18(3): 893-898. <https://doi.org/10.13057/biodiv/d180305>
- Madduppa H., Timm, J. & Kochzius, M. 2014. Interspecific, spatial and temporal variability of self-recruitment in Anemonefishes. *Plos One*. 9(2): 1-13. <https://doi.org/10.1371/journal.pone.0090648>
- Madduppa, H.H., Bahri, S., Subhan, B., Anggraini, N.P., Ohoiulun, H., Abdillah, T., Arafat, D., Santoso, P. & Sangadji, I.M. 2019. DNA barcoding of sea turtles (Dermochelyidae and Cheloniidae) and its protocol using different tissues quality: Implication to conservation managers. *IOP Conf. Ser. Earth Environ. Sci.*, 278(1): 012041. <https://doi.org/10.1088/1755-1315/278/1/012041>
- Manuputty A.E.W. 1996. Soft coral is one of the constituents of coral reefs. *Oceana*. 11 (4):131-141 (in Bahasa Indonesia)
- Manuputty A.E.W. 2010. Distribution of soft coral, genus *Sinularia* May, 1898 (Octocorallia, Alcyonacea) on Derawan islands, East Kalimantan. *Oceanologi dan Limnologi di Indonesia*. 36: 211-225. (in Indonesian)
- Manuputty A.E.W. & van Ofwegen, L.P. 2007. The genus *Sinularia* (Octocorallia: Alcyonacea) from Ambon and Seram (Moluccas, Indonesia). *Zool. Med. Leiden*. 81: 187-216.
- Manuputty AEW. 2002, Soft coral Indonesian waters, LIPI Pusat Penelitian Oseanografi. Jakarta. 91 pp
- Mantelatto, M.C., da Silva, A.G., dos Santos Louzada, T., McFadden, C.S. & Creed, J.C., 2018. Invasion of aquarium origin soft corals on a tropical rocky reef in the southwest Atlantic, Brazil. *Mar. Poll. Bull.*, 130: 84-89. <https://doi.org/10.1016/j.marpolbul.2018.03.014>
- McFadden, C.S., France, S.C., Sánchez, J.A. & Alderslade, P., 2006a. A molecular phylogenetic analysis of the Octocorallia (Cnidaria: Anthozoa) based on mitochondrial protein-coding sequences. *Mol. Phylogenet. Evol.*, 41: 513-527. <https://doi.org/10.1016/j.ympev.2006.06.010>
- McFadden, C.S., Alderslade, P., Van Ofwegen, L.P., Johnsen, H. & Rusmevichientong, A. 2006b. Phylogenetic relationships within the tropical soft coral genera *Sarcophyton* and *Lobophytum* (Anthozoa, Octocorallia). *Invertebr. Biol.*, 125: 288-305. <https://doi.org/10.1111/j.1744-7410.2006.00070.x>
- McFadden C.S. & van Ofwegen. L.P. 2012. *Stoloniferous octocorals* (Anthozoa, Octocorallia) from South Africa, with descriptions of a new family of Alcyonacea, a new genus of Clavulariidae and a new species of Cornularia (Cornulariidae). *Invertebr. Syst.* 26: 331-356. <https://doi.org/10.1071/IS12035>
- McFadden, C.S., Van Ofwegen, L.P., Beckman, E.J., Benayahu, Y. & Alderslade, P.. 2009. Molecular systematics of the speciose Indo-Pacific soft coral genus, *Sinularia* (Anthozoa: Octocorallia). *Invertebr. Biol.*, 128: 303-323. <https://doi.org/10.1111/j.1744-7410.2009.00179.x>
- McFaddrn, C.S., Benayahu, Y., Pante, E., Thoma, J.N., Nevarez, P.A. & France, S.C.,. Limitations of mitochondrial gene barcoding in Octocorallia. *Mol. Ecol. Res.*, 11: 19-31. <https://doi.org/10.1111/j.1755-0998.2010.02875.x>
- McFadden, C.S., Reynolds, A.M. & Janes, M.P., 2014. DNA barcoding of xeniid soft corals (Octocorallia: Alcyonacea: Xeniidae) from Indonesia: Species richness and phylogenetic relationships. *System. Biodivers.*, 12: 247-257. <https://doi.org/10.1111/syb.12035>

org/10.1080/14772000.2014.902866

- Moore K.M., Alderslade, P. & Miller, K.J. 2017. A taxonomic revision of Anthothela (Octocorallia: Scleraxonia: Anthothelidae) and related genera, with the addition of new taxa, using morphological and molecular data. *Zootaxa*, 4304(1): 1-212. <https://doi.org/10.11646/zootaxa.4304.1.1>
- Prehadi, P., Sembiring, A., Kurniasih, E.M., Rahmad, R., Arafat, D., Subhan, B. & Madduppa, H.H., 2015. DNA barcoding and phylogenetic reconstruction of shark species landed in Muncar fisheries landing site in comparison with Southern Java fishing port. *Biodiversitas*. 16: 55-61. doi: 10.13057/biodiv/d160107
- Puillandre, N., Lambert, A., Brouillet, S. & Achaz, G.J.M.E., 2012. ABGD, Automatic Barcode Gap 543 Discovery for primary species delimitation. *Mol. Ecol.* 21: 1864-544. <https://doi.org/10.1111/j.1365-294X.2011.05239.x>
- Quattrini, A.M., Etnoyer, P.J., Doughty, C., English, L., Falco, R., Remon, N., Rittinghouse, M. & Cordes, E.E., 2014. A phylogenetic approach to octocoral community structure in the deep Gulf of Mexico. *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 99: 92-102. <https://doi.org/10.1016/j.dsr2.2013.05.027>
- Rao, C.R. 1964. The use and interpretation of principal component analysis in applied research. *Sankhyā*, 26: 329-359.
- Riegl B., Schleyer, M.H., Cook, P.J. & Branch, G.M. 1995. Structure of Africa's southernmost coral communities. *Bull. Mar. Sci.* 56: 676-691.
- Roxas, H.A. 1933. Philippine Alcyonaria. The families Cornulariidae and Xenidae. *Philippine J. Sci.* 50: 49-110
- Saleky D., Setyobudiandi, I., Toha, H.A., Takdir, M. & Madduppa, H. 2016. Length-weight relationship and population genetic of two marine gastropods species (Turbinidae: Turbo sparverius and Turbo bruneus) in the Bird Seascape Papua, Indonesia. *Biodiversitas*. 17: 208-217. <https://doi.org/10.13057/biodiv/d170130>
- Sanger F., Nicklen, S. & Coulson, A.R. 1977. DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci.* 74(12): 5463-5467. <https://doi.org/10.1073/pnas.74.12.5463>
- Schenk A., 1896. Clavulariiden, Xenidiiden und Alcyoniiden von Ternate. *Abhandl. Senckenb. Naturf. Ges.*, Ed. 23: 41-80. <https://doi.org/10.5962/bhl.title.9164>
- Sembiring, A., Pertiwi, N.P.D., Mahardini, A., Wulandari, R., Kurniasih, E.M., Kuncoro, A.W., Cahyani, N.D., Anggoro, A.W., Ulfa, M., Madduppa, H. & Carpenter, K.E., 2015. DNA barcoding reveals targeted fisheries for endangered sharks in Indonesia. *Fish. Res.* 164: 130-134. <https://doi.org/10.1016/j.fishres.2014.11.003>
- Sheu, J.H., Peng, B.R., Fang, L.S., Hwang, T.L., Su, J.H., Wu, Y.C. & Sung, P.J. 2019 Hydroperoxyditerpenoids from Octocorals. *Isr. J. Chem.* 59: 403-413. <https://doi.org/10.1002/ijch.201900011>.
- Tamura, K., Stecher, G., Peterson, D., Filipksi, A. & Kumar, S. 2013. Mega 5: Molecular Evolution Genetics Analysis Version 6.0. *Mol. Biol. Evo.* 30: 2725-2729. <https://doi.org/10.1093/molbev/mst197>
- Toha, A.H.A., Dailami, M., Anwar, S., Setiawan, J.B., Jentewo, Y., Lapadi, I., Sutanto, S, Aryasari, R., Ambariyanto, A., Runtuboi, F. & Madduppa, H. 2020. The genetic relationships and Indo-Pacific connectivity of whale sharks (*Rhincodon typus*) with particular reference to mitochondrial COI gene sequences from Cendrawasih Bay, Papua, Indonesia. *Biodiversitas*, 21(5): 2159-2171. <https://doi.org/10.13057/biodiv/d210544>
- Tursch, B. & Tursch, A. 1982. The soft coral community on a sheltered reef quadrat at Laing Island (Papua New Guinea). *Mar. Biol.* 68: 321-332. <https://doi.org/10.1007/BF00409597>
- Utinomi, H. 1953. Invertebrate fauna of the intertidal zone of the Tokara Islands. VI. Alcyonaria. *Pub. Seto Mar. Biol. Lab.* 3(2): 149-160, <https://doi.org/10.5134/174473>
- Vogler, C., Benzie, J.A.H., Tenggardjaja, K., Ambariyanto, A., Barber, P.H. and Wörheide, G., 2013. Phylogeography of the crown-of-thorns starfish: genetic structure within the Pacific species. *Coral Reefs*, 32(2): 515-525.
- Veron, J.E., Devantier, L.M., Turak, E., Green, A.L.,

Kininmonth, S., Stafford-Smith, M. & Peterson, N. 2009. Delineating the coral triangle. *Galaxea*, 11: 91-100. <https://doi.org/10.3755/galaxea.11.91>

Wabnitz C., Taylor, M., Green, E. & Razak, T. 2003. *From Ocean to Aquarium*. UNEP-WCMC, Cambridge, UK. 72 pp