The future of the past: Applications of paleoecological findings in peatland restoration in Indonesia

K. Anggi Hapsari¹, T.C. Jennerjahn^{2,3} and H. Behling¹

There are several potential practical applications of paleoecological information that can provide guidance for improving peatland restoration in Indonesia, and highlight the values, roles, and applicability of paleoecology in ecological restoration.

The importance of peatlands in Indonesia Indonesian peatlands provide habitat for a large diversity of organisms, including endangered flora and fauna. They also store ~56 Gt carbon, equivalent to five years of current global carbon emissions, mainly in their soggy, spongy soil layers made of partially decomposed dead plant material. In the past few decades, rapid peatland degradation and drainage (Fig. 1) has not only posed a threat to the native orangutans and other species living there, who are rapidly losing their habitat, but has also facilitated peat fires, leading to haze and related health crises, and costing Indonesia billions of dollars in economic loss (World Bank 2019). Peat fires also released enormous amounts of carbon dioxide, fueling climate warming, and placing Indonesia amongst the world's top carbon emitters. It is not surprising there are growing calls for restoration.

"Ecological restoration can be improved using paleoecology": a statement easily found in many paleo-related publications. However, there is a dearth of examples on



Figure 1: Images of (A) pristine, (B) degraded, and (C) burnt peatlands in Indonesia (photo credits: A. Hapsari and I. Fikri).

how paleo-information can be practically applied in restoration beyond defining prehuman-intervention baselines. Meanwhile, attempting to restore the ecosystems to their pre-human conditions is increasingly being considered to be unattainable or even inappropriate (Clewell and Aronson 2013; Baker and Eckerberg 2016; Rull, this issue). Here, we use a paleorecord from the Sungai Buluh peatland in Indonesia as our main example, where the impact of the Malayu Empire's activities and how the peatland responded to them, is documented (Fig. 2; Hapsari et al. 2018; 2021).

Goal adjustment

When defining what it is we are trying to restore, we can use ecological attributes of a successfully restored or undegraded ecosystem (e.g. species composition, resilience, functions) as goals (Clewell and Aronson 2013). We then need to ensure the goals are realistic and can be met within regulatory constraints and available resources.

Most, if not all, peatland restoration projects in Indonesia aim to restore both hydrology and peat swamp forests to an intact state, i.e. prior to disturbance, and refurbish their functions and values (Puspitaloka et al. 2020). These projects are typically planned for 60 years, but many are shorter (3-30 years) due to limited resources.

The paleorecord from Sungai Buluh taught us that it took 170 years for the peatland to return to its pre-disturbance condition after "light" human use (i.e. no clearing/drainage). Thus, the goals of present-day peatland restoration are guite unrealistic, as we now face much harder ecological (drained/cleared/ burnt) and socioeconomic (resistance/sabotage) challenges, a much shorter timeframe, and often limited resources. It is necessary for the projects to either opt for easier goals (e.g. zero-illegal logging, blockage of all canals) or to extend the project durations, especially if the fulfillment/unfulfillment of the set goals will decide the project's fate, e.g. further funding, permit extension, or penalty.

Refine the designs

The Sungai Buluh record suggests that a peatland can fully recover if it isn't drained and still has some of the old forest remaining. Restoration projects for peatlands with these conditions can choose the low-cost

prescribed natural regeneration (passive restoration) approach. The resources available can be concentrated to remove/reduce the disturbances, e.g. fire prevention, law enforcement, ecoliteracy and bioeconomy development, or alternative livelihood creation.

For the drained and deforested peatlands, assisted natural regeneration or partial/ complete reconstruction is needed, which includes replanting and/or rewetting. However, although replantation in peatlands creates an alternative livelihood for the locals, it is very expensive and often fails because unsuitable taxa are used, e.g. dryland species are planted in deeply flooded conditions (canal/burnt scar). A paleorecord from Borneo shows that peatlands sometimes developed in Pandanus-dominated freshwater lakes (Hope et al. 2005). Replantation can then mimic natural succession by planting Pandanus in the deeply flooded areas (lake-like conditions) and introduce other desirable taxa once those areas are shallow enough (Giesen and Sari 2018).

For restoration to be successful, the target ecosystem needs to eventually manage to self-organize, an attribute that is mainly driven by the planted taxa's regenerating ability. The recovery of the Sungai Buluh peat swamp forest started with recruitment of taxa with high seed productivities, viabilities, germination rates, and seedling survival rates. Peatland restoration plans thus need to include those and/or other taxa with similar regenerative power and, if necessary, to introduce specific pollinators or seed dispersers (e.g. *Trigona* bees for *Shorea*).

Paleorecords also suggest that dense forest cover is the secret of peatlands' fire-resistance (Hapsari et al. 2021). To improve target peatlands' resilience to fire, forest cover needs to be grown quickly. While the current strategy chooses to plant in single-species groups, introducing a few individuals from many species and letting the "selection effect" do the work is a better option than planting a large number of poorly adapted species and having none survive.

Stream of income

In restoration, socioeconomic achievement is important, but it won't be sustainable long-term without ecological success and durability. Subsequent management of



restored peatland needs to ensure that the socioeconomic activities won't jeopardize peatland resilience. The Sungai Buluh record suggests that collection of non-timber forest products and fodder, and selective/restrictive wood extraction without draining the land or burning the trees, are sustainable. A paleorecord from Borneo (Hope et al. 2005) also shows that human impact was restricted to the riversides because of access difficulties. Restoration can mimic the low level of past human disturbance by concentrating economically valuable taxa alongside natural waterways to ease people's access to their income source, thus limiting the temptation to reopen blocked canals, which cause major challenges in peatland rewetting.

Time is of the essence

Restoration, unfortunately, isn't a quick fix, but a long-term effort that requires patience and dedication (Clewell and Aronson 2013). Policy in Indonesia constrains the durations of peatland restoration projects within 60 (extendable to 100) years. Assuming that target peatlands will also require about 170 years to recover, like the Sungai Buluh, the 60-year framework is far too short. In fact, the target peatlands are often drained, thus requiring physical environment modification (rewetting), which will likely prolong the recovery time.

A major driver for restoration success is the time that has elapsed since the restoration started (Crouzeilles et al. 2017). Thus, the current policy needs to allow for some flexibility in granting permissible periods according to the projects' goals, e.g. longer duration for projects targeting a full biodiversity recovery than for projects aiming at zero-carbon emission (Hapsari et al. 2018). If the current policy remains in place, it needs to, at least, consider the substantial time lag between the completion of project tasks and the attainment of restoration targets. The 60(-100)-year permissible project length can serve as an "intervention period"; however, the restoration site should remain protected and monitored for another 60(-100) years to accommodate the development of peatlands' self-organization and self-sustainability.

A hope... and a warning!

Many people wonder whether ecological restoration is really achievable, or if it is just empty promises and a massive waste of investment (Conniff 2015; Almassi 2017). As the success of these efforts remains to be seen, the long wait may give rise to growing scepticism (Giesen and Sari 2018; Miller et al. 2021). The documented recovery of the Sungai Buluh peatland from past human disturbance certainly gives hope for tropical peatland restoration efforts as it shows that recovery is a slow process and that decades of waiting do not necessarily mean failure. However, it should also be taken as a cautionary example that it will take a whole lot longer to depave a parking lot and build back a paradise than the other way aroundas in Joni Mitchell's song.*

*Big Yellow Taxi: jonimitchell.com/music/song. cfm?id=13



Figure 2: The Sungai Buluh peatland underwent major changes throughout the Malayu Empire occupation (orange shade; modified from Hapsari et al. 2018; 2021): (A) Peat swamp forest cover declined soon after the Malayu Empire arrival (~1050 cal yr BP); (B) A lower proportion of high-quality timber trees and historically recorded natural resource use for construction by the Malayu Empire suggests selective timber collection; (C) Vigorous growth of unpalatable forage that is usually driven by the lack of competition from other palatable herbs suggests fodder collection; (D) An increase in Piperaceae while the Malayu Empire became the largest pepper center in Sumatra implies non-timber forest product (NTPF) collection; (E) Relatively low charcoal concentration and peak magnitude suggests inextensive fire use; (F) Continuously accumulating peat substrate that needs a consistently high water-table suggests an absence of land drainage; (G) PSF regeneration after ME abandonment started with the increase of rapidly regenerating taxa.

AFFILIATIONS

- ¹Department of Palynology and Climate Dynamics, University of Göttingen, Germany
- ²Department of Biogeochemistry and Geology, Leibniz Centre for Tropical Marine Research (ZMT),
- Bremen, Germany
- ³Faculty of Geoscience, University of Bremen, Germany

CONTACT

K. Anggi Hapsari: kartika.hapsari@biologie.unigoettingen.de

REFERENCES

Almassi B (2017) Ethics Environ 22: 19-40

Baker S, Eckerberg K (2016) Restor Ecol 24: 284-290 Clewell AF, Aronson J (2013) Ecological restoration: principles, values, and structure of an emerging profession. Island Press, 303 pp

- Conniff R (2015) The false promise of ecological restoration projects. TakePart Magazine
- Crouzeilles R et al. (2017) Sci Adv 3: e1701345
- Giesen W, Sari ENN (2018) Tropical Peatland Restoration Report: The Indonesian Case. Euroconsult Mott MacDonald, 98 pp
- Hapsari KA et al. (2018) J Ecol 106: 1-18
- Hapsari KA et al. (2021) Rev Palaeobot and Palynol 293: 104482
- Hope G et al. (2005) Quat Res 64: 407-417
- Miller MA et al. (2021) Sustain Dev 1-15
- Puspitaloka D et al. (2020) Restor Ecol 28: 435-446
- World Bank (2019) The World Bank Annual Report 2019: Ending Poverty, Investing in Opportunity. World Bank, 95 pp

PA()ES